

## GENERAL INFORMATION

**Cyto-Pak™ Micro** contains micronutrients that have been complexed with chelating agents. It should be applied at planting or at the beginning of stressful growth stages such as first true leaf, flowering, and fruit set. **Cyto-Pak™ Micro** improves efficiency of photosynthesis, increases the rate of CO<sub>2</sub> fixation, and enhances the plant's ability to use available nutrients. **Cyto-Pak™ Micro** is processed with organic compounds, including enzymes, hydrolyzed proteins, and amino acids, that are the end result of microbial fermentation.

**Cyto-Pak™ Micro** is recommended for use on row crops and vegetables.

## GUARANTEED ANALYSIS

### Secondary and Minor Elements

Copper (Cu).....0.10%  
(Derived from micronized copper sulfate.)

## COMPATIBILITY

Do not mix this product with conventional 2,4-D amine products as a compatibility problem may occur.

If mixing with fertilizers, it is recommended that a small scale test for compatibility be conducted before mixing a full batch.

Information regarding the contents and levels of metals in this product is available on the internet at:


<http://www.aapfco.org/metals.htm>

Net contents: 5 gallons

Weight per gallon: 9.88 lbs.

# CYTO-PAK™ MICRO

A complexed Micronutrient Package  
with Organic Compounds  
for Soil and Foliar Application



P.O. Box 3453 Pasco WA 99302

(509) 547-8234

**SHAKE WELL BEFORE USING**

**HARMFUL IF SWALLOWED**

## DIRECTIONS FOR USE AND APPLICATION RATES

### Soil:

1. At the time of planting, apply 8 - 16 oz./acre **Cyto-Pak™ Micro** with starter fertilizer, banded or dribbled in the seed furrow, or in irrigation water. Alternatively, apply 8 - 16 oz./acre **Cyto-Pak™ Micro** incorporated with fertilizer at the first side-dress.

(Washington State law requires that EPA-approved safety devices be in use when this product is used in any irrigation system.)

2. At the second irrigation or side-dress, apply 8 - 16 oz./acre **Cyto-Pak™ Micro**. This product may be tank-mixed with fertilizers.

### Foliar:

1. Three weeks after emergence, apply 4 - 8 oz./acre **Cyto-Pak™ Micro**. This product may be tank-mixed with pesticides and fertilizers.

2. Three or four weeks after the first application, apply 4 - 8 oz./acre **Cyto-Pak™ Micro**.

3. Seven to ten days after the second application, apply 4 - 8 oz./acre **Cyto-Pak™ Micro**.

### CONDITION OF SALE

Seller warrants that the product conforms to its chemical description and is reasonable fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. Since weather, crop, soil, and other conditions may vary, Northwest Ag Products, Inc. and/or seller make no warranty of any kind, expressed or implied, concerning the use of this product. The user assumes all risk of use of handling, whether or not in accordance with directions or suggestions.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**